

# GALOIS MODULE STRUCTURE OF MILNOR K-THEORY MOD $p^s$ IN CHARACTERISTIC $p$

JÁN MINÁČ\*, ANDREW SCHULTZ, AND JOHN SWALLOW†

ABSTRACT. Let  $E$  be a cyclic extension of  $p$ th-power degree of a field  $F$  of characteristic  $p$ . For all  $m, s \in \mathbb{N}$ , we determine  $K_mE/p^s K_mE$  as a  $(\mathbb{Z}/p^s\mathbb{Z})[\text{Gal}(E/F)]$ -module. We also provide examples of extensions for which all of the possible nonzero summands in the decomposition are indeed nonzero.

Let  $F$  be a field of characteristic  $p$ . Let  $K_m F$  denote the  $m$ th Milnor  $K$ -group of  $F$  and  $k_m F = K_m F/pK_m F$ . (See, for instance, [Mi] and [Ma, Chapter 14].) If  $E/F$  is a Galois extension of fields, let  $G = \text{Gal}(E/F)$  denote the associated Galois group. In [BLMS] the structure of  $k_mE$  as an  $\mathbb{F}_p G$ -module was determined when  $G$  is cyclic of  $p$ th-power order. In this paper we determine the Galois module structure of  $K_mE$  modulo  $p^s$  for  $s \in \mathbb{N}$  and these same  $G$ . We also provide examples of extensions for which the possible free summands in the decomposition are all nonzero. These examples together with the results in [BLMS] show that the dimensions over  $\mathbb{F}_p$  of indecomposable  $\mathbb{F}_p[\text{Gal}(E/F)]$ -modules occurring as direct summands of  $k_mE$  are all powers of  $p$  and that all dimensions  $p^i$ ,  $i = 0, 1, \dots, n$ , indeed occur in suitable examples.

Recall the theorem of Bloch-Kato and Gabber (see [BK]): the sequence

$$0 \rightarrow k_m F \rightarrow \Omega_F^m \xrightarrow{\mathfrak{P}} \Omega_F^m/d\Omega_F^{m-1}$$

is exact, where  $\Omega_F^m$  is the  $m$ th graded component of the exterior algebra on Kähler differentials and  $\mathfrak{P}$  is the Artin-Schreier operator. In [I, §6], Izhboldin succeeded in providing an analogue of this important

---

*Date:* September 5, 2007.

*2000 Mathematics Subject Classification.* 19D45, 12F10.

*Key words and phrases.* Milnor  $K$ -theory, cyclic extension, Galois module, norm.

\*Research supported in part by NSERC grant R0370A01, and by a Distinguished Research Professorship at the University of Western Ontario.

†Research supported in part by NSF grant DMS-0600122.

interpretation of  $k_m F$ , as follows: for  $s \in \mathbb{N}$ , the sequence

$$0 \rightarrow K_m F / p^s K_m F \xrightarrow{\delta} Q^m(F, s) \xrightarrow{\mathfrak{P}} Q^m(F, s)$$

is exact, where  $Q^m(F, s)$  is the “Milnor-Witt” group of  $F$ , defined together with maps  $\delta$  and  $\mathfrak{P}$  in [I, §6]. These objects play an important role in the arithmetic of fields, higher class field theory, and Milnor  $K$ -theory of fields of characteristic  $p$ . (See, for instance, [FK] and the references therein.)

Since the classification problem of  $(\mathbb{Z}/p^s\mathbb{Z})G$ -modules for cyclic  $G$  is nontrivial and has not been completely solved—see, for instance, [T] for results and references—it is a pleasant surprise that the  $(\mathbb{Z}/p^s\mathbb{Z})G$ -modules  $K_m E / p^s K_m E$  have a simple description. The main ingredients we use to obtain this description are the lack of  $p$ -torsion in  $K_m E$ , due to Izhboldin [I], together with the result [BLMS] for the case  $s = 1$ —which also depends on Izhboldin’s result.

Suppose that  $E/F$  is cyclic of degree  $p^n$ , and for  $i = 0, 1, \dots, n$ , let  $E_i/F$  be the subextension of degree  $p^i$  of  $E/F$  and  $G_i := \text{Gal}(E_i/F)$ . Set  $R_s := \mathbb{Z}/p^s\mathbb{Z}$  and let  $\mathbb{Z}_p$  be the ring of  $p$ -adic integers. We write  $\iota_{F,E}: K_m F \rightarrow K_m E$  and  $N_{E/F}: K_m E \rightarrow K_m F$  for the natural inclusion and norm maps, and we use the same notation for the induced maps between  $K_m F / p^s K_m F$  and  $K_m E / p^s K_m E$ .

**Theorem.** *There exists an isomorphism of  $R_s G$ -modules*

$$K_m E / p^s K_m E \simeq \bigoplus_{i=0}^n Y_i$$

where

- $Y_n$  is a free  $R_s G$ -module of rank  $\dim_{\mathbb{F}_p} N_{E/F} k_m E$ ,
- $Y_i$ ,  $0 < i < n$ , is a free  $R_s G_i$ -module of rank

$$\dim_{\mathbb{F}_p} N_{E_i/F} k_m E_i / N_{E_{i+1}/F} k_m E_{i+1},$$

- $Y_0$  is a free  $R_s$ -module of rank

$$\dim_{\mathbb{F}_p} k_m F / N_{E_1/F} k_m E_1 \text{ with } Y_0^G = Y_0,$$

and

- for each  $0 \leq i \leq n$ ,  $Y_i \subseteq \iota_{E_i,E}(K_m E_i / p^s K_m E_i)$ .

Moreover,

$$\widehat{K_m E} := \varprojlim_s K_m E / p^s K_m E \simeq \bigoplus_{i=0}^n \hat{Y}_i,$$

where each  $\hat{Y}_i$  is a free  $\mathbb{Z}_p G_i$ -module of the same rank as  $Y_i$ .

## 1. PROOF OF THE THEOREM

We prove the result by induction on  $s$ . The case  $s = 1$  is [BLMS, Theorem 2]. Assume therefore that  $s > 1$  and the result holds for  $s - 1$ :

$$K_mE/p^{s-1}K_mE = \oplus \tilde{Y}_i,$$

with each  $\tilde{Y}_i$  a free  $R_{s-1}G_i$ -module  $\tilde{Y}_i$  in the image of  $\iota_{E_i, E}$ ,  $0 \leq i \leq n$ .

For each  $i$  with  $0 \leq i \leq n$ , let  $B_{s-1,i} \subset \iota_{E_i, E}K_mE_i/p^{s-1}K_mE_i$  be an  $R_{s-1}G_i$ -base for the free  $R_{s-1}G_i$ -module  $\tilde{Y}_i$ . By induction the cardinality of  $B_{s-1,i}$  is

$$|B_{s-1,i}| = \begin{cases} \dim_{\mathbb{F}_p} N_{E_i/F}k_mE_i/N_{E_{i+1}/F}k_mE_{i+1}, & i < n \\ \dim_{\mathbb{F}_p} N_{E/F}k_mE, & i = n. \end{cases}$$

Since the  $\tilde{Y}_i$  are independent, the set

$$B_{s-1} := \cup_{0 \leq i \leq n} B_{s-1,i} \subseteq K_mE/p^{s-1}K_mE$$

is  $R_{s-1}G$ -independent.

For each  $i$ , let  $\mathcal{B}_i \subseteq \iota_{E_i, E}(K_mE_i)$  be a set of representatives for the elements of  $B_{s-1,i}$ , and let  $B_{s,i} \subseteq K_mE/p^sK_mE$  be chosen to make the following first diagram commutative. The second diagram merely recalls where our  $\mathcal{B}_i$ ,  $B_{s,i}$  and  $B_{s-1,i}$  are located.

$$\begin{array}{ccc} \mathcal{B}_i & \xrightarrow{\text{mod } p^s} & B_{s,i} \\ \text{mod } p^{s-1} \downarrow & \swarrow \text{mod } p^{s-1} & \\ B_{s-1,i} & & \end{array} \quad \begin{array}{ccc} K_mE & \longrightarrow & K_mE/p^sK_mE \\ \downarrow & \swarrow & \\ K_mE/p^{s-1}K_mE & & \end{array}$$

Hence for each  $i$  we have bijections

$$\mathcal{B}_i \leftrightarrow B_{s,i} \leftrightarrow B_{s-1,i}$$

and  $|B_{s,i}| = |B_{s-1,i}|$ .

First we observe that every nonzero ideal  $V$  of  $R_sG_i$  contains  $p^{s-1}(\tau - 1)^{p^i}$ , where  $\tau$  is any fixed generator of  $G_i$ . Indeed consider  $0 \neq \beta \in B$ . By multiplying by an appropriate power of  $p$ , we may assume  $0 \neq \beta \in p^{s-1}R_sG_i$ . Let us write

$$\beta = \sum_{j=k}^{p^i-1} c_j(\tau - 1)^j,$$

where each  $c_j \in p^{s-1}R_s$ ,  $j = k, \dots, p^i - 1$ , and  $c_k \notin p^sR_s = \{0\}$ , say  $c_k = p^{s-1}\tilde{c}_k$  with  $\tilde{c}_k \notin pR_s$ . Using the fact that  $p^{s-1}(\tau - 1)^{p^i} = 0$  in  $R_sG_i$  we see that we can multiply  $\beta$  by  $(\tau - 1)^{p^i-k-1}$  to obtain

$$0 \neq \tilde{c}_k p^{s-1}(\tau - 1)^{p^i-1} \in V.$$

Since  $\tilde{c}_k \in U(R_s)$  we see that  $p^{s-1}(\tau - 1)^{p^i-1} \in V$  as asserted.

Set  $Y_i$  to be the  $R_sG$ -submodule of  $K_mE/p^sK_mE$  generated by  $B_{s,i}$ . It is clear that  $Y_i \subseteq \iota_{E_i,E}(K_mE_i/p^sK_mE_i)$  and hence  $Y_i$  is an  $R_sG_i$ -module.

Each element  $b \in B_{s,i}$  generates in  $K_mE/p^sK_mE$  a free  $R_sG_i$ -module  $M_b$ , as follows. Suppose that  $M_b$  is not a free  $R_sG_i$ -module. Then the annihilator of  $b$  in  $R_sG_i$  is a nonzero ideal of  $R_sG_i$ . Let  $\hat{b} \in \mathcal{B}_i$  and  $\tilde{b} \in B_{s-1,i}$  correspond to  $b$  under the bijection above. Let also  $\sigma$  be a generator of  $G$  and  $\bar{\sigma}$  its image in  $G_i$ . Since every nonzero ideal of  $R_sG_i$  contains  $p^{s-1}(\bar{\sigma} - 1)^{p^i-1}$ , for some  $\hat{c} \in K_mE$  we have

$$p^{s-1}(\bar{\sigma} - 1)^{p^i-1}\hat{b} = p^s\hat{c}.$$

Since  $K_mE$  has no  $p$ -torsion [I, Theorem A], we obtain

$$p^{s-2}(\bar{\sigma} - 1)^{p^i-1}\hat{b} = p^{s-1}\hat{c}.$$

Then in  $K_mE/p^{s-1}K_mE$

$$p^{s-2}(\bar{\sigma} - 1)^{p^i-1}\tilde{b} = 0,$$

contradicting the fact that  $\tilde{b}$  lies in the  $R_{s-1}G_i$ -base  $B_{s-1,i}$  for  $\tilde{Y}_i$ . (Alternatively we could use [T, Theorem 5.1] to show that  $M_b$  is a free  $R_sG_i$ -module.)

Now set  $B_s := \cup_{0 \leq i \leq n} B_{s,i} \subseteq K_mE/p^sK_mE$ . Suppose we have a relation

$$\sum r_\alpha b_\alpha = 0, \quad b_\alpha \in B_s, \quad r_\alpha \in R_sG_\alpha,$$

such that  $G_\alpha = G_i$  for a suitable  $i$ ,  $0 \leq i \leq n$ , with  $b_\alpha \in B_{s,i}$ . Let  $\tilde{b}_\alpha \in B_{s-1}$  correspond to  $b_\alpha$  under the natural bijection, and similarly let  $\tilde{r}_\alpha \in R_{s-1}G_\alpha$  be the image of  $r_\alpha \in R_sG_\alpha$ .

Working mod  $p^{s-1}$  we have  $\sum \tilde{r}_\alpha \tilde{b}_\alpha = 0$ . Since each  $\tilde{b}_\alpha$  lies in the  $R_{s-1}G$ -independent set  $B_{s-1}$ , we deduce that  $r_\alpha \in p^{s-1}R_sG_\alpha$  for each  $\alpha$ . Write  $r_\alpha = ps_\alpha$  for elements  $s_\alpha \in R_sG_\alpha$ . We rewrite the original relation as

$$\sum ps_\alpha b_\alpha = 0, \quad b_\alpha \in B_s, \quad s_\alpha \in R_sG_\alpha.$$

Just as before we divide by  $p$  to obtain in  $K_m E / p^{s-1} K_m E$

$$\sum \tilde{s}_\alpha \tilde{b}_\alpha = 0.$$

Again since each  $\tilde{b}_\alpha \in B_{s-1}$ , we deduce that  $s_\alpha \in p^{s-1} R_s G_\alpha$ . But then  $r_\alpha = p s_\alpha = 0 \in R_s G_\alpha$  for each  $r_\alpha$ , as desired.

Hence for each  $i$  in  $0 \leq i \leq n$  we have that  $Y_i$  is a direct sum of free  $R_s G_i$ -modules  $M_b$  for  $b \in B_{s,i}$ , and moreover that  $\sum Y_i = \oplus Y_i$ . By Nakayama's Lemma, since  $\mathcal{B}$  generates  $K_m / p^{s-1} K_m E$  it also generates  $K_m / p^s K_m E$ , and hence  $\oplus Y_i = K_m E / p^s K_m E$ . (More explicitly, choose  $\alpha \in K_m E / p^s K_m E$  and  $\hat{\alpha} \in K_m E$  a lift of  $\alpha$ . Since  $B_{s-1}$  spans  $K_m E / p^{s-1} K_m E$  we have

$$\hat{\alpha} = \sum_{b \in \mathcal{B}} f_b b + p^{s-1} \hat{\gamma}$$

for some  $\hat{\gamma} \in K_m E$ , where each  $f_b \in R_s G$  and all but finitely many  $f_b = 0$ . We also have  $\hat{\gamma} = \sum g_b b + p^{s-1} \hat{\delta}$  for some  $\hat{\delta} \in K_m E$ , where again each  $g_b \in R_s G$  and all but finitely many  $g_b = 0$ . Therefore

$$\alpha = \sum_{b \in \mathcal{B}} (f_b + p^{s-1} g_b) b$$

as elements of  $K_m E / p^s K_m E$ .)

The last statement concerning the equality  $\widehat{K_m E} = \oplus_{i=0}^n \hat{Y}_i$  is obtained by passing to projective limits.  $\square$

## 2. EXAMPLES OF $E/F$ WITH $Y_i \neq \{0\}$ FOR ALL $i$

Let  $p$  be an arbitrary prime number, and let  $q$  be an arbitrary prime number or 0. We show that for each  $n, m \in \mathbb{N}$  there exists a cyclic field extension  $E/F$  of degree  $p^n$  and characteristic  $q$  such that for each  $i$ ,  $0 \leq i < n$ ,

$$\dim_{\mathbb{F}_p} N_{E_i/F} k_m E_i / N_{E_{i+1}/F} k_m E_{i+1} \neq 0$$

and

$$\dim_{\mathbb{F}_p} N_{E/F} k_m E \neq 0.$$

Recall that we index  $E_i$  such that  $F \subset E_i \subset E$  and  $[E_i : F] = p^i$ .

We are interested in the images of the quotient groups in  $k_m E$ . Because in the case  $p = q$  the natural homomorphism  $k_m F \rightarrow k_m E$  is injective, for the case of our main application we can work in  $k_m F$ .

### 2.1. The case $m = 1$ .

Fix  $p$ ,  $q$ , and  $n$  as above and set  $m = 1$ . We construct a field extension  $E/F$  as above together with elements

$$\begin{aligned} x_i &\in N_{E_i/F}(E_i^\times) \setminus N_{E_{i+1}/F}(E_{i+1}^\times)F^{\times p}, \quad 0 \leq i < n, \\ x_n &\in N_{E/F}(E^\times) \setminus F^{\times p}. \end{aligned}$$

Let  $B$  be a field of characteristic  $q$ , and let  $A/B$  be a cyclic extension of degree  $p^n$ . Index the subfields  $A_i$  of  $A/B$  such that  $[A_i : B] = p^i$ , and denote by  $\iota_{B,A_i} : K_1 B \hookrightarrow K_1 A_i$  the natural inclusion. Let  $\sigma$  be a generator of  $\text{Gal}(A/B)$  and set  $\sigma_i = \sigma|_{A_i}$ , the restricted map. Finally, assume that there exist elements  $x_0, x_1, \dots, x_n \in B^\times$  such that the following condition  $(*)$  holds:

$$\begin{aligned} [\iota_{B,A_j}(x_j)]^{p^{n-j-1}} &\notin \langle [\iota_{B,A_j}(x_1)]^{p^{n-1}}, [\iota_{B,A_j}(x_2)]^{p^{n-2}}, \dots, [\iota_{B,A_j}(x_n)] \rangle \\ &\subseteq A_j^\times / N_{A/A_j}(A^\times), \quad 0 \leq j < n, \\ x_n &\notin B^{\times p}, \end{aligned}$$

where  $[x]$  denotes the class of  $x$  and  $\langle S \rangle$  the subgroup generated by a set  $S$  in the named factor group. At the end of this section we shall create an example where condition  $(*)$  holds.

Now consider cyclic algebras

$$\mathcal{A}_j = (A/B, \sigma, x_j^{p^{n-j}}), \quad 1 \leq j \leq n.$$

Observe that

$$[\mathcal{A}_j] = [(A_j/B, \sigma_j, x_j)] \in \text{Br}(B), \quad 1 \leq j \leq n$$

([P, Chapter 15, Corollary b]), where  $\text{Br}(B)$  denotes the Brauer group of  $B$ . Let  $F$  be the function field of the product of Brauer-Severi varieties of  $\mathcal{A}_1, \dots, \mathcal{A}_n$ . (See [SV, page 735]; see also [J, Chapter 3] for basic properties of Brauer-Severi varieties.)

Let  $E$  be the compositum  $A \cdot F$  of the fields  $F$  and  $A$ . Since  $F$  is a regular extension of  $B$ , we see that  $E/F$  is a cyclic extension of degree  $p^n$ . We denote again as  $\sigma$  the generator of  $\text{Gal}(E/F)$  which restricts to  $\sigma \in \text{Gal}(A/B)$ , and we write  $E_k = A_k \cdot F$  for  $k = 0, 1, \dots, n$ . Now  $[\mathcal{A}_j \otimes_B F] = 0 \in \text{Br}(F)$ ,  $j = 1, \dots, n$ , because  $F$  splits each  $\mathcal{A}_j$ . Hence

$$0 = [(E/F, \sigma, x_j^{p^{n-j}})] = [(E_j/F, \sigma_j, x_j)],$$

and so  $x_j \in N_{E_j/F}(E_j^\times)$  as desired (see [P, Chapter 15, page 278]).

However, we claim that

$$\begin{aligned} x_j &\notin (N_{E_{j+1}/F}(E_{j+1}^\times))F^{\times p}, \quad 0 \leq j < n, \\ x_n &\notin F^{\times p}. \end{aligned}$$

Since  $x_n \notin B^{\times p}$  by hypothesis and  $F/B$  is a regular extension, we have  $x_n \notin F^{\times p}$ . Assume then that  $0 \leq j < n$  and, contrary to our statement,

$$x_j \in (N_{E_{j+1}/F}(E_{j+1}^\times))F^{\times p}.$$

Then we have  $x_j f^p \in N_{E_{j+1}/F}(E_{j+1}^\times)$  for some  $f \in F^\times$ . Hence

$$[(E_{j+1}/F, \sigma_{j+1}, x_j f^p)] = 0 \in \text{Br}(F)$$

and so

$$\begin{aligned} [(E_{j+1}/F, \sigma_{j+1}, x_j)] &= -[(E_{j+1}/F, \sigma_{j+1}, f^p)] \\ &= -[(E_j/F, \sigma_j, f)]. \end{aligned}$$

(In the case  $j = 0$ , we use  $(E_0/F, \sigma_0, f)$  to denote the zero element in  $\text{Br}(F)$ .) Consequently  $(E_{j+1}/F, \sigma_{j+1}, x_j)$  is split by  $E_j$ . (See [P, Chapter 15, Proposition b].)

But then

$$[(E_{j+1}/E_j, \sigma_{j+1}^{p^j}, \iota_{F, E_j}(x_j))] = 0 \in \text{Br}(E_j).$$

(See [D, page 74].) Hence  $[(E/E_j, \sigma^{p^j}, \iota_{F, E_j}(x_j^{p^{n-j-1}}))] = 0 \in \text{Br}(E_j)$ . But  $E_j = A_j \cdot F$  is the function field of the product of the Brauer-Severi varieties of  $\mathcal{A}_k \otimes_B A_j$  for  $k = 1, \dots, n$ . Therefore

$$[(A/A_j, \sigma^{p^j}, \iota_{B, A_j}(x_j^{p^{n-j-1}}))] \in \langle [\mathcal{A}_k \otimes_B A_j], k = 1, \dots, n \rangle \subseteq \text{Br}(A_j).$$

Consequently

$$\begin{aligned} [\iota_{B, A_j}(x_j^{p^{n-j-1}})] &\in \langle [\iota_{B, A_j}(x_1)]^{p^{n-1}}, \dots, [\iota_{B, A_j}(x_n)] \rangle \\ &\in A_j^\times / N_{A/A_j}(A^\times), \end{aligned}$$

a contradiction to condition (\*).

Thus we have shown that a required extension  $E/F$  exists with elements  $x_0, x_1, \dots, x_n$ , provided that we can produce a field extension  $A/B$  and elements  $x_0, x_1, \dots, x_n \in B^\times$  such that condition (\*) is valid. Now we show that such an extension and elements exist.

Let  $B := C(x_0, x_1, \dots, x_n)$ , where  $C$  is a field with characteristic  $q$  and  $x_0, x_1, \dots, x_n$  are algebraically independent elements over  $C$ . Assume also that there exists a cyclic extension  $D/C$  of degree  $p^n$  with

Galois group  $G = \langle \sigma \rangle$ . Finally, let  $A := D(x_0, x_1, \dots, x_n)$ . Thus  $A/B$  is a cyclic extension of degree  $p^n$ .

We claim that condition  $(*)$  holds. Clearly  $x_n \notin B^{\times p}$ . Contrary to our claim assume that

$$x_j^{p^{n-j-1}} = x_1^{c_1 p^{n-1}} \cdots x_n^{c_n} N_{A/A_j}(\gamma)$$

where  $c_i \in \mathbb{Z}$ ,  $\gamma \in A^\times$ , and  $0 \leq j < n$ . Write

$$\gamma = u \cdot \frac{p_1 \cdots p_t}{q_1 \cdots q_w},$$

where  $\{p_1, \dots, p_t\}$  and  $\{q_1, \dots, q_w\}$  are disjoint sets of primes in the ring  $D[x_0, x_1, \dots, x_n]$  and  $u \in U(D[x_0, x_1, \dots, x_n]) = D^\times$ .

We write  $H_j = \text{Gal}(A/A_j)$  for  $j = 0, 1, \dots, n-1$ . Then we have

$$x_j^{p^{n-j-1}} \prod_{h \in H_j} \prod_{i=1}^w h(q_i) = x_1^{c_1 p^{n-1}} \cdots x_n^{c_n} N_{A/A_j}(u) \prod_{h \in H_j} \prod_{i=1}^t h(p_i).$$

In order to show that this equation is impossible, consider the discrete valuation  $v_j$  on  $A$  such that  $v_j(x_j) = 1$ ,  $v_j(x_k) = 0$  if  $j \neq k$ , and  $v_j(d) = 0$  for  $d \in D$ . Consider first the case when  $x_j \nmid q_l$ ,  $l = 1, \dots, w$ . Then the value of  $v_j$  on the left-hand side is  $p^{n-j-1}$ , while the value of  $v_j$  on the right-hand side is at least  $p^{n-j}$ . Indeed if  $c_j \neq 0$  this is true as  $x_j^{p^{n-j}}$  divides the right-hand side of our equation. If  $c_j = 0$  or  $j = 0$  (in which case  $c_0$  is not defined) then since  $x_j$  divides the left-hand side we see that there exists  $p_l$ ,  $l \in \{1, \dots, t\}$ , such that  $x_j \mid p_l$ . Because  $|H_j| = p^{n-j}$  we see again that the value of  $v_j$  on the right-hand side is again at least  $p^{n-j}$ . Thus  $x_j \mid q_l$  for some  $l = 1, \dots, w$ , and so  $x_j \nmid p_l$  for any  $l = 1, \dots, t$ . Then the value of  $v_j$  on the left-hand side of our equation is  $p^{n-j-1} + p^{n-j}$  while the value of  $v_j$  on the right-hand side is at most  $p^{n-j}$ . Thus we see that the equation above is impossible. Hence condition  $(*)$  is valid and we have established the desired example of  $E/F$  in the case  $m = 1$ .

## 2.2. The case $m > 1$ .

Fix  $m$ ,  $n$ , and  $q$  as above and let  $L/K$  be a field extension satisfying the case  $m = 1$  with the same  $n$  and  $q$ . Let  $x_0, x_1, \dots, x_n \in K^\times$  such that

$$\begin{aligned} x_i &\in N_{L_i/K}(L_i^\times) \setminus N_{L_{i+1}/K}(L_{i+1}^\times) K^{\times p}, \quad 0 \leq i < n, \\ x_n &\in N_{L/K}(L^\times) \setminus K^{\times p}. \end{aligned}$$



Consider the field of the iterated power series  $F := K((y_1)) \cdots ((y_{m-1}))$ . Then  $E := L \cdot F$  is a cyclic extension of degree  $p^n$  over  $F$ . For each  $j \in \{0, 1, \dots, n\}$  consider the element

$$\alpha_j = \{x_j, y_1, \dots, y_{m-1}\} \in k_m F.$$

(If  $m = 1$  then  $\alpha_j = \{x_j\}$ .) By our hypothesis and the projection formula for the norm map in  $K$ -theory, we have

$$\alpha_j \in N_{E_j/F} k_m E_j.$$

Now for  $0 \leq j < n$  we shall prove by induction on  $m \in \mathbb{N}$  that

$$\alpha_j \notin N_{E_{j+1}/F} k_m E_{j+1}.$$

If  $m = 1$  then our statement is true by the choice of the field extension  $L/K$  and the elements  $x_j$ . Assume then that  $m > 1$  and that our statement is true for  $m - 1$ .

Consider the complete discrete valuation  $v$  on  $F$  with uniformizer  $y_{m-1}$  and residue field  $F_v = K((y_1)) \cdots ((y_{m-2}))$  if  $m > 2$  and  $F_v = K$  if  $m = 2$ .

For the sake of simplicity we denote by  $E'$  the field  $E_{j+1}$ , and denote the unique extension of  $v$  on  $E'$  again by  $v$ . Since we are considering an unramified extension we assume that both valuations are normalized. Let  $\partial : k_m F \rightarrow k_{m-1} F_v$  and  $k_m E' \rightarrow k_{m-1} E'_v$  be the homomorphisms induced by residue maps in Milnor  $K$ -theory. Then applying [K, Lemma 3] we see that the following diagram is commutative:

$$\begin{array}{ccc} k_m E' & \xrightarrow{\partial} & k_{m-1} E'_v \\ N_{E'/F} \downarrow & & \downarrow N_{E'_v/F_v} \\ k_m F & \xrightarrow{\partial} & k_{m-1} F_v. \end{array}$$

If  $\alpha_j \in N_{E'/F} k_m E'$ , then  $\partial \alpha_j \in N_{E'_v/F_v} k_{m-1} E'_v$ .

But  $\partial \alpha_j = \{x_j, y_1, \dots, y_{m-2}\}$  if  $m > 2$  and  $\partial \alpha_j = \{x_j\}$  if  $m = 2$ , a contradiction in either case. Therefore we have constructed a field extension  $E/F$  with the desired properties.

**Remarks.** In [MSS1] we determined the  $\mathbb{F}_p G$ -module structure of  $k_1 E$  for all cyclic extensions  $E/F$  of degree  $p^n$ , where  $G$  is the Galois group. In particular, the decomposition does not depend upon the characteristic of the base field. The ranks of the free  $\mathbb{F}_p G_i$ -summands appearing in that decomposition are again determined by the image of  $N_{E_i/F}(E_i^\times)/N_{E_{i+1}/F}(E_{i+1}^\times)$  in  $E^\times/E^{\times p}$ .

When no primitive  $p$ th root of unity lies in  $F$ , we have that  $F^\times/F^{\times p}$  embeds in  $E^\times/E^{\times p}$ . Therefore the construction given above for field extensions  $E/F$  applies in this case, and these free  $\mathbb{F}_p G_i$ -summands do indeed occur for any characteristic. When a primitive  $p$ th root of unity is in  $F^\times$ , the kernel of the homomorphism  $F^\times/F^{\times p} \rightarrow E^\times/E^{\times p}$  is generated by a class  $[a] \in F^\times/F^{\times p}$ , where  $E_1 = F(\sqrt[p]{a})$ . In this case it is enough to additionally require that  $a \in N_{E/F}(E^\times)$  when  $p = 2$  and  $n = 1$  (since the condition is automatic otherwise), and the construction above of field extensions  $E/F$  applies again.

If  $F$  contains a primitive  $p$ th-root of unity and  $m = 1$ , the decomposition contains at most one other indecomposable module, a cyclic  $\mathbb{F}_p G$ -module of dimension  $p^k + 1$  for  $k \in \{-\infty, 0, 1, \dots, n-1\}$  (where we set  $p^{-\infty} = 0$ ). In [MSS2] we showed that all of these modules are realizable as well.

#### ACKNOWLEDGMENTS

Andrew Schultz would like to thank Ravi Vakil for his encouragement and direction in this and in all other projects. We would also like to thank Ganesh Bhandari for his useful comments related to this paper.

#### REFERENCES

- [BK] S. Bloch and K. Kato.  $p$ -adic étale cohomology. *Inst. Hautes Études Sci. Publ. Math.* No. 63 (1986), 107–152.
- [BLMS] G. Bhandari, N. Lemire, J. Mináč, and J. Swallow. Galois module structure of Milnor  $K$ -theory in characteristic  $p$ . Preprint arXiv:math.NT/0405503v5 (2007).
- [D] P. K. Draxl. *Skew fields*. London Mathematical Society Lecture Note Series 81. Cambridge: Cambridge University Press, 1983.
- [FK] I. Fesenko and M. Kurihara, eds. *Invitation to higher local fields (Münster, 1999)*. Geometry & Topology Monographs 3. Somerville, MA: International Press, 2000.
- [I] O. Izhboldin. On  $p$ -torsion in  $K_*^M$  for fields of characteristic  $p$ . *Algebraic K-theory*. Advances in Soviet Mathematics 4. Providence, RI: American Mathematical Society, 1991, 129–144.
- [J] N. Jacobson. *Finite-dimensional division algebras*. New York: Springer-Verlag, 1996.
- [K] K. Kato. Residue homomorphisms in Milnor  $K$ -theory. *Advanced Studies in Pure Math.* **2** (1983), 153–172.
- [Ma] B. Magurn. An algebraic introduction to  $K$ -theory. *Encyclopedia of Mathematics and its Applications* 87. Cambridge: Cambridge University Press, 2002.

- [Mi] J. Milnor. Algebraic  $K$ -theory and quadratic forms. *Invent. Math.* **9** (1970), 318–344.
- [MSS1] J. Mináč, A. Schultz, and J. Swallow. Galois module structure of  $p$ th-power classes of cyclic extensions of degree  $p^n$ . *Proc. London Math. Soc.* **92** (2006), no. 2, 307–341.
- [MSS2] J. Mináč, A. Schultz, and J. Swallow. Cyclic algebras and construction of some Galois modules. Preprint arXiv:math.NT/0410536v1 (2004).
- [P] R. Pierce. *Associative algebras*. Graduate Texts in Mathematics 88. New York: Springer-Verlag, 1982.
- [SV] A. Schofield and M. Van den Bergh. The index of a Brauer class on a Brauer-Severi variety. *Trans. Amer. Math. Soc.* **333** (1992), no. 2, 729–739.
- [T] J. Thévenaz. Representations of finite groups in characteristic  $p^r$ . *J. Algebra* **72** (1981), 478–500.

DEPARTMENT OF MATHEMATICS, MIDDLESEX COLLEGE, UNIVERSITY OF  
WESTERN ONTARIO, LONDON, ONTARIO N6A 5B7 CANADA

*E-mail address:* minac@uwo.ca

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF ILLINOIS AT URBANA-CHAM-  
PAIGN, 273 ALTGELD HALL, MC-382, 1409 W. GREEN STREET, URBANA, IL  
61801 USA

*E-mail address:* acs@math.uiuc.edu

DEPARTMENT OF MATHEMATICS, DAVIDSON COLLEGE, BOX 7046, DAVIDSON,  
NORTH CAROLINA 28035-7046 USA

*E-mail address:* joswallow@davidson.edu